

Product Datasheet GenieFluor 488 Anti-Mouse CD3ε Antibody [145-2C11] Catalogue Code: AGEL0377

Antibody Data

Product SKU:	AGEL0377	Clone:	145-2C11
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	T-cell surface glycoprotein CD3 epsilon chain;CD3E;T-cell surface antigen T3/Leu-4 epsilon chain;CD3e;CD3E;T3E; P22646 CD3ɛ is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the lg superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3ɛ forms a TCR complex by associating with the CD3δ, γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100	
Size:	50 Tests, 100 Tests, 200 Tests	80 - S	
Host Species:	Armenian Hamster	(%) 60	
Isotype:	Armenian Hamster IgG	$\mathbf{E}_{2}^{40} \underbrace{1_{0}^{400}}_{350} \underbrace{1_{0}^{400}}_{400} \underbrace{1_{50}^{400}}_{500} \underbrace{1_{50}^{500}}_{550} \underbrace{1_{000}^{500}}_{600} \underbrace{1_{000}^{500}}_{650} \underbrace{1_{000}^{700}}_{Wavelength (nm)}$	
Isotype Control:	-		

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.