

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL0582</b>	<b>Clone:</b>	<b>2.4G2</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** CD16a/b;CD32;CD32A/B;Fc fragment of IgG low affinity IIa/b receptor;Fc fragment of IgG low affinity IIIa/b receptor;Fc fragment of IgG low affinity IIIb receptor;Fc gamma receptor III A/B;FCG2A;FcGR;FCGR2A/BFCGR3;FCGR3A/B;Fc gamma RIIa/b ;

**Uniprot ID:** P08508 P08101

**Background:** CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and mediate adaptive immune responses.


**Form:** Liquid

**Conjugation:** PE/Cyanine 7

**Size:** 50 Tests, 100 Tests, 200 Tests

**Host Species:** Rat

**Isotype:** Rat IgG2b, κ



Ex:495;565;755 nm; Em:775 nm

**Isotype Control:** PE/Cyanine7 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL0582]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

---