

#### **Product Datasheet**

# PerCP/Cyanine5.5 Anti-Mouse CD3 Antibody [17A2]

Catalogue Code: AGEL0662

# Antibody Data

Product SKU: AGEL0662 Clone: 17A2

Applications: FCM

Reactivity: Mouse

### **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: T-cell surface glycoprotein CD 3epsilon/delta/gamma/zeta

chain;CD3E/D/G/Z;CD3e/d/g/z;CD3E/D/G/Z;CD3;

**Uniprot ID:** P04235 P11942 P22646 P24161

**Background**: CD3, also known as T3, is a member of the Ig superfamily and primarily expressed on T

cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3 $\epsilon$ ,  $\delta$ ,  $\gamma$  and  $\zeta$  chains. It forms a TCR complex by associating with TCR  $\alpha/\beta$  or  $\gamma/\delta$  chains. CD3 plays a critical role in TCR signal transduction, T cell activation,

and antigen recognition by binding the peptide/MHC antigen complex.

Form: Liquid


**Conjugation:** PerCP/Cyanine 5.5

Size: 25µg, 100µg

Host Species: Rat

**Isotype:** Rat IgG2b, κ

PerCP/Cyanine5.5 Excitation and Emission Spectra



Ex:440;480;675 nm; Em:675 nm

**Isotype Control:** PerCP/Cyanine5.5 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL0662]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/106 cells in 100  $\mu$ L

volume].