

Product Datasheet GenieFluor 647 Anti-Mouse CD11a Antibody [FD441.8] Catalogue Code: AGEL0839

Antibody Data

Product SKU:	AGEL0839	Clone:	FD441.8	
Applications:	FCM			
Reactivity:	Mouse			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	Integrin alpha-L;Itgal;CD11 antigen-like family member A;LFA-1A;Lymphocyte antigen 15;Ly-15;CD11a; P24063 CD11a is a 180 kD glycoprotein, also known as αL integrin, LFA-1 α, Ly-15, or Ly-21. It is a member of the integrin family, primarily expressed on lymphocytes, monocytes/macrophages, and granulocytes. In association with CD18, the CD11a/CD18 complex forms LFA-1. CD11a plays an important role in intercellular adhesion and costimulation by binding its ligands, ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).		
Form:	Liquid	647 Excitation and Emission Spectra	
Conjugation:	Genie Fluor647	100	
Size:	25µg, 100µg	80 - S	
Host Species:	Rat	(%) 100 - 00 - 00 - 00 - 00 - 00 - 00 - 00	
Isotype:	Rat IgG2b, κ	$\mathbf{F}_{20} = \begin{pmatrix} 40 & & & \\ 20 & & & \\ 0 & & & \\ 350 & 400 & 450 & 500 & 550 & 600 & 650 & 700 & 750 & 800 & 850 \\ \hline \mathbf{Wavelength} (nm) \end{pmatrix}$	

Isotype Control: Genie Fluor 647 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL0839]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].