

Product Datasheet **PE/Cyanine5.5 Anti-Human CD41 Antibody [HIP8]** Catalogue Code: AGEL1136

Antibody Data

Product SKU:	AGEL1136	Clone:	HIP8	
Applications:	FCM			
Reactivity:	Human			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	Integrin alpha-lib;ITGA2B;GPalph lib;CD41;GP2B; ITGAB; P08514	na lib;GPIIb;Platelet membrane glycoprotein	
Background:	CD41 is a 125/25 kD α subunit of the gpIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (gpIIb α) and light chain (gpIIb β) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor and thrombin.		
Form:	Liquid	PE/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5.5	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	€ 60	
Isotype:	Mouse IgG1, κ	2 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:495;565;675 nm; Em:690 nm	

Isotype Control: PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1136]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.