

Product Datasheet GenieFluor 488 Anti-Mouse CD3ɛ

Antibody [145-2C11] Catalogue Code: AGEL1329

Antibody Data

Product SKU:	AGEL1329	Clone:	145-2C11
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	T-cell surface glycoprotein CD3 epsilon chain;CD3E;T-cell surface antigen T3/Leu-4 epsilon chain;CD3e;CD3E;T3E; P22646		
Background:	CD3 ϵ is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 ϵ forms a TCR complex by associating with the CD3 δ , γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100	
Size:	25µg, 100µg	80	
Host Species:	Armenian Hamster		
Isotype:	Armenian Hamster IgG	$\frac{1}{2}$ $\frac{1}{400}$ $\frac{1}{400}$ $\frac{1}{450}$ $\frac{1}{500}$ $\frac{1}{500}$ $\frac{1}{500}$ $\frac{1}{600}$ $\frac{1}{650}$ $\frac{1}{700}$ Wavelength (nm)	

Isotype Control: Genie Fluor 488 Armenian Hamster IgG Isotype Control[PIP] [Product AGEL1329]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].