

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL1369</b>	<b>Clone:</b>	<b>AFS98</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

**Important Note:**


Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** Macrophage colony-stimulating factor 1 receptor;CSF-1 receptor (EC:2.7.10.1);CSF-1-R;CSF-1R;M-CSF-R;Proto-oncogene c-Fms;CD115;Csf1r;Csfmr;Fms;  
**Uniprot ID:** P09581

**Background:** CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have described an Ig-like extracellular domain, a transmembrane domain, an intracellular juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

<b>Form:</b>	Liquid
<b>Conjugation:</b>	APC
<b>Size:</b>	25µg, 100µg
<b>Host Species:</b>	Rat
<b>Isotype:</b>	Rat IgG2a, κ



<b>Isotype Control:</b>	APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL1369]
<b>Storage Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10<sup>6</sup> cells in 100 µL volume].