

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL1437</b>	<b>Clone:</b>	<b>GR-20</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		


**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** Interferon gamma receptor 1;Ifngr1;IFN-gamma-R1;IFN-gamma-R-alpha;CD119;  
**Uniprot ID:** P15261  
**Background:** CDw119 is a 90 kD immunoglobulin superfamily member, also known as IFN- $\gamma$ R $\alpha$  chain. It is a class II cytokine receptor family member that serves as a IFN- $\gamma$ -binding chain associated with the IFN- $\gamma$   $\beta$  chain also known as AF-1. In addition to ligand binding, CDw119 participates in ligand trafficking. CDw119 is expressed on T and B cells, NK cells, fibroblasts, endothelial, and epithelial cells. Binding of IFN- $\gamma$  induces receptor dimerization, internalization, Jak1 and Jak2 protein kinase activation and, ultimately, STAT1 activation. IFN- $\gamma$  initiates and regulates a variety of immune responses.

**Form:** Liquid  
**Conjugation:** PE  
**Size:** 50 Tests, 100 Tests, 200 Tests  
**Host Species:** Rat  
**Isotype:** Rat IgG2a,  $\kappa$



Ex:495;565 nm; Em:575 nm

**Isotype Control:** PE Rat IgG2a,  $\kappa$  Isotype Control[2A3] [Product AGEL1437]  
**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.  
**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

---