

Product Datasheet **FITC Anti-Mouse CD45.2 Antibody [104.2]** Catalogue Code: AGEL1525

Antibody Data

Applications: FCM	Product SKU:	AGEL1525	Clone:	104.2	
	Applications:	FCM			
Reactivity: Mouse	Reactivity:	Mouse			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Ly-5.2; LCA;		
Uniprot ID:	-		
Background:	CD45.2 is an alloantigen of CD45, expressed by Ly5.2 bearing mouse strains (e.g., A, AKR, BALB/c, CBA/Ca, CBA/J, C3H/He, C57BL, C57BR, C57L, C58, DBA/1, DBA/2, NZB, SWR, 129). CD45, a member of the protein tyrosine phosphatase (PTP) family, is a 180-240 kD glycoprotein expressed on all hematopoietic cells except mature erythrocytes and platelets. There are multiple isoforms in the mouse that play key roles in TCR and BCR signal transduction. These isoforms are very specific to the activation and maturation states of the cell as well as specific cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	100	
Size:	25µg, 100µg		
Host Species:	Mouse	60 - 40 - 40 - 40 - 40 - 40 - 40 - 40 -	
Isotype:	Mouse IgG2a, к	20 0 350 400 450 500 550 600 650 700 Wavelength (nm) Ex:490 nm; Em:530 nm	

Isotype Control: FITC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL1525]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].