

#### **Product Datasheet**

# GenieFluor 647 Anti-Mouse CD279/PD-1 Antibody [29F.1A12]

Catalogue Code: AGEL1631

### **Antibody Data**

Product SKU: AGEL1631 Clone: 29F.1A12

Applications: FCM

Reactivity: Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information:**

**Alternate Names:** PD-1; Programmed Death-1;

Uniprot ID: Q02242

**Background**: CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging

to the CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative

costimulation is essential for prolonged survival of intratesticular islet allografts.


Form: Liquid

**Conjugation:** Genie Fluor647

Size: 50 Tests, 100 Tests, 200 Tests

Host Species: Rat

**Isotype:** Rat IgG2a, κ



**Isotype Control:** Genie Fluor 647 Rat IgG2a, κ Isotype Control[2A3] [Product AGEL1631]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



Stability & Storage:

Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.