

#### **Product Datasheet**

# Purified Anti-Mouse CD274/PD-L1 Antibody [10F.9G2]

Catalogue Code: AGEL1637

## Antibody Data

Product SKU: AGEL1637 Clone: 10F.9G2

Applications: FCM

Reactivity: Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information:**

Alternate Names: B7-H1; PD-L1; Programmed cell death ligand 1;B7 homolog 1;B7-H; B7H1; PDL1;

PDCD1L1; PDCD1LG1;

Uniprot ID: Q9EP73

Background: CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I

transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. It is expressed on T cells, B cells, NK cells, dendritic cells, IFN- $\gamma$  activated endothelial cells, and monocytes. B7-H1 is one of the ligands of PD-1. The interaction of B7-H1 with PD-1 plays an important role in the inhibition of T cell responses. Other studies have shown that B7-H1 is able to costimulate T cell growth and cytokine production. CD274 is involved in costimulation essential for T cell proliferation and production of IL-10 and IFN- $\gamma$ , in an IL-2-dependent and a PD-1-independent manner. Its interaction with PD-

1 inhibits T cell proliferation and cytokine production.

Form: Liquid

Conjugation: Unconjugated

Size: 25µg, 100µg

Host Species: Rat

**Isotype:** Rat IgG2b, κ

**Isotype Control:** Purified Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL1637]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq$  1.0 µg per 106 cells in 100 µL volume or 100 µL of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.