

Product Datasheet

GenieFluor 488 Anti-Human CD10 Antibody [HI10a] Catalogue Code: AGEL1721

## Antibody Data

Product SKU:	AGEL1721	Clone:	HI10a	
Applications:	FCM			
Reactivity:	Human			

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID:	Neprilysin;Atriopeptidase;CALLA;Enkephalinase;NEP;Neutral endopeptidase;Skin fibroblast elastase;SFE;CD10; P08473		
Background:	CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins, bradykinin, angiotensins I and II, and other biologically active peptides.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100	
Size:	20 Tests, 100 Tests, 200 Tests	80 - S	
Host Species:	Mouse	NA 10	
Isotype:	<b>уре:</b> Mouse IgG1, к	E 40 20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	

Isotype Control: Genie Fluor 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1721]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.