

Product Datasheet **PE/Cyanine5.5 Anti-Human CD90 Antibody [5E10]** Catalogue Code: AGEL1833

## Antibody Data

Product SKU:	AGEL1833	Clone:	5E10
Applications:	FCM		
Reactivity:	Human		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	CDw90;T25;FLJ33325;Thy1;		
Uniprot ID:	P04216		
Background:	CD90 is a 25-35 kD GPI-anchored protein, also known as Thy-1. It belongs to the Ig superfamily. Human CD90 is expressed on neuronal cells, a subset of CD34+ cells, a subset of fetal liver cells and fetal thymocytes, fibroblasts, activated endothelial cells, and some leukemia cell lines. CD34+CD90+ cells are primitive hematopoietic stem cells. It has been reported that Thy-1 binds with $\beta$ 2 and $\beta$ 3 integrins and plays bimodal roles in the regulation of cell adhesion and neurite outgrowth, and inhibits hematopoietic stem cells proliferation and differentiation.		
Form:	Liquid	PE/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5.5	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	(%) 500 100 100 100 100 100 100 100 100 100	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (mm)	
		En 105-5(5/75 mm En /00 mm	

Ex:495;565;675 nm; Em:690 nm

**Isotype Control:** PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1833]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.