

Product Datasheet **APC Anti-Mouse CD24 Antibody [M1/69]** Catalogue Code: AGEL1893

## Antibody Data

Product SKU:	AGEL1893	Clone:	M1/69
Applications:	FCM		
Reactivity:	Mouse		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID: Background:	Cd24a; Ly-52;HAS;Nectadrin;R13-Ag;X62 heat stable antigen ; P24807 CD24 is a 35-45 kD protein also known as Heat Stable Antigen (HSA), Ly-52, or Nectadrin. It is a GPI-linked sialoglycoprotein expressed on lymphocytes, granulocytes, epithelial cells, thymocytes, monocytes, erythrocytes, and dendritic cells. CD24 expression varies during T and B cell differentiation and is a useful marker for delineating various lymphocyte developmental stages. CD24 serves as an adhesion or costimulatory molecule involved in T and B lymphocyte activation and differentiation by homophilic binding or binding to CD62P.		
Form:	Liquid	APC Excitation and Emission Spectra	
Conjugation:	APC	100 -	
Size:	50 Tests, 100 Tests, 200 Tests		
Host Species:	Rat	(\$) - 00 - 00 - 00 - 00 - 00 - 00 - 00 - 0	
Isotype:	Rat IgG2b, κ	20 0 350 400 450 500 550 550 600 650 700 750 800 850 850 Wavelength (nm) Ex:650 nm; Em:660 nm	

**Isotype Control:** APC Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL1893]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.