

Product Datasheet

GenieFluor 647 Anti-Mouse CD24 Antibody [M1/69] Catalogue Code: AGEL1908

Antibody Data

Product SKU:	AGEL1908	Clone:	M1/69	
Applications:	FCM			
Reactivity:	Mouse			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Cd24a; Ly-52;HAS;Nectadrin;R13-Ag;X62 heat stable antigen ;		
Uniprot ID:	P24807	(nown on Linet Ctable Antigen (LICA), Ly, 50, er Negtedrig	
Background:	CD24 is a 35-45 kD protein also known as Heat Stable Antigen (HSA), Ly-52, or Nectadrin. It is a GPI-linked sialoglycoprotein expressed on lymphocytes, granulocytes, epithelial cells, thymocytes, monocytes, erythrocytes, and dendritic cells. CD24 expression varies during T and B cell differentiation and is a useful marker for delineating various lymphocyte developmental stages. CD24 serves as an adhesion or costimulatory molecule involved in T and B lymphocyte activation and differentiation by homophilic binding or binding to CD62P.		
Form:	Liquid	647 Excitation and Emission Spectra	
Conjugation:	Genie Fluor647	100 -	
Size:	25µg, 100µg	80 ·	
Host Species:	Rat	(%) 60 . 40 .	
Isotype:	Rat IgG2b, κ	20 0 350 400 450 550 550 600 650 700 750 800 850 Wavelength (nm)	

Isotype Control: Genie Fluor 647 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL1908]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].