

Product Datasheet

APC Anti-Mouse FcεRlα Antibody [MAR-

Catalogue Code: AGEL1986

Antibody Data

Product SKU: AGEL1986 Clone: MAR-1

Applications: FCM

Reactivity: Mouse

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: High affinity immunoglobulin epsilon receptor subunit alpha;Fc-epsilon RI-

alpha;FcERI;Fcer1a;

Uniprot ID: P20489

Background: Fc ϵ Rl α is a transmembrane protein belonging to the Ig superfamily. Fc ϵ Rl α forms a

tetrameric complex with one β and two γ -subunits. The Fc ϵ RI complex plays an important role in triggering IgE-mediated allergic reactions. It is abundantly expressed on mast and basophils and up-regulated by the presence of IgE. Following stimulation via Fc ϵ RI α , mast cells and basophils release bioactive chemical mediators such as histamine, resulting in the initiation of allergic reactions. Cross linking of the high-affinity receptor for IgE on tissue mast cells triggers immediate hypersensitivity with local symptoms. The MAR-1

monoclonal antibody reacts with the FcεRlα subunit.


Form: Liquid

Conjugation: APC

Size: 25µg, 100µg

Host Species: Armenian Hamster

Isotype: Armenian Hamster IgG



Ex:650 nm; Em:660 nm

Isotype Control: APC Armenian Hamster IgG Isotype Control[PIP] [Product AGEL1986]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/106 cells in 100 μ L

volume].