

Product Datasheet

PE/Cyanine5.5 Anti-Mouse/Human CD11b Antibody [M1/70] Catalogue Code: AGEL2169

## Antibody Data

Applications: FCM	
Reactivity: Human;Mouse	

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID:	Integrin alpha-M;Itgam;CD11 antiger adhesion receptor MO1;CD11b; P05555 P11215	n-like family member B;CR-3 alpha chain;Leukocyte	
Background:	CD11b is a 170 kD glycoprotein also known as $\alpha$ M integrin, Mac-1 $\alpha$ subunit, Mol, CR3, and Ly-40. CD11b is a member of the integrin family, primarily expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b non-covalently associates with CD18 ( $\beta$ 2 integrin) to form Mac-1. Mac-1 plays an important role in cell-cell interaction by binding its ligands ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen.		
Form:	Liquid	PE/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5.5	100	
Size:	50 Tests, 100 Tests, 200 Tests		
Host Species:	Rat		
Isotype:	Rat IgG2b, κ	Z 20 0 350 400 450 550 550 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:495;565;675 nm; Em:690 nm	

Isotype Control: PE/Cyanine5.5 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL2169]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.