

#### **Product Datasheet**

# GenieFluor 488 Anti-Mouse CD117 Antibody [2B8]

Catalogue Code: AGEL2181

# Antibody Data

Product SKU: AGEL2181 Clone: 2B8

Applications: FCM

Reactivity: Mouse

### **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Mast/stem cell growth factor receptor Kit;Kit;SCFR;Proto-oncogene c-Kit;Tyrosine-protein

kinase Kit;CD117;

Uniprot ID: P05532

**Background**: CD117 is a 145 kD immunoglobulin superfamily member also known as c-Kit and stem

cell factor receptor (SCFR). It is a transmembrane tyrosine-kinase receptor that binds the c-Kit ligand (also known as steel factor, stem cell factor, and mast cell growth factor). CD117 is expressed on hematopoietic stem cells (including multipotent hematopoietic stem cells, progenitors committed to myeloid and/or erythroid lineages, and T and B cell precursors), mast cells, and acute myeloid leukemia (AML) cells. CD117 interaction with

its ligand is critical for the development of hematopoietic stem cells.


Form: Liquid

**Conjugation:** Genie Fluor488

Size: 25µg, 100µg

Host Species: Rat

**Isotype:** Rat IgG2b, κ



**Isotype Control:** Genie Fluor 488 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL2181]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/106 cells in 100  $\mu$ L volume].