

Product Datasheet

Low Endotoxin Purified Anti-Mouse CD90.1 Antibody [19E12] Catalogue Code: AGEL2189

## Antibody Data

| Product SKU:  | AGEL2189 | Clone: | 19E12 |
|---------------|----------|--------|-------|
| Applications: | FCM      |        |       |
| Reactivity:   | Mouse    |        |       |
|               |          |        |       |

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

| Alternate Names: | Thy-1.1 membrane glycoprotein;Thy1.1;Thy-1.1 antigen;CD90.1;Thy-1.1;   |  |
|------------------|--|--|
| Uniprot ID:      | -  |  |
| Background:      | CD90.1 is a 25-35 kD immunoglobulin superfamily member, also known as Thy-1. It is expressed on hematopoietic stem cells and neurons, all thymocytes, and peripheral T cells. CD90.1 is a glycosylphosphatidylinositol (GPI)-anchored membrane glycoprotein involved in signal transduction. CD90.1 is involved in costimulation of lymphocyte proliferation and activation, and hematopoietic stem cell differentiation. CD90.1 has been shown to interact with CD45. |  |
| Form:            | Liquid   |  |
| Conjugation:     | None (AF/LE)   |  |
| Size:            | 50µg, 500µg, 1mg   |  |
| Host Species:    | Rat  |  |
| Isotype:         | Rat IgG2a, к   |  |
|                  |  |  |

| Isotype Control: | AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product AGEL2189]   |
|------------------|---|
| Storage Buffer:  | 0.2 μm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay. |
| Shipping:        | Biological ice pack at 4°C  |



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \ \mu$ g per 106 cells in 100  $\mu$ L volume or 100  $\mu$ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.