

Product Datasheet PE Anti-Mouse CD45R/B220 Antibody [RA3.3A 1/6.1] Catalogue Code: AGEL2215

Antibody Data

| Product SKU: | AGEL2215 | Clone: | RA3.3A 1/6.1 |
|---------------|----------|--------|--------------|
| Applications: | FCM | | |
| Reactivity: | Mouse | | |
| | | | |

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

| Alternate Names: | B220; | | |
|------------------|---|--|--|
| Uniprot ID: | - | | |
| Background: | CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-Fas ^{lpr} and MRL-Fas ^{gld} mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity. | | |
| Form: | Liquid | PE Excitation and Emission Spectra | |
| Conjugation: | PE | 100 - | |
| Size: | 25µg, 100µg | | |
| Host Species: | Rat | € 60 - Titer 40 - | |
| Isotype: | Rat IgM, κ | 20 0 350 400 450 500 550 600 650 700 Wavelength (nm) | |
| | | Ex:495;565 nm; Em:575 nm | |
| Isotype Control: | PE Rat IgM, κ Isotype Control[RTK2118] [Product AGEL2215] | | |
| Storage Buffer: | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. | | |

Storage Buffer:

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].