

#### **Product Datasheet**

# FITC Anti-Mouse FcεRIα Antibody [MAR-

Catalogue Code: AGEL2314

### Antibody Data

Product SKU: AGEL2314 Clone: MAR-1

Applications: FCM

Reactivity: Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information:**

Alternate Names: High affinity immunoglobulin epsilon receptor subunit alpha;Fc-epsilon RI-

alpha;FcERI;Fcer1a;

Uniprot ID: P20489

**Background**: Fc $\epsilon$ Rl $\alpha$  is a transmembrane protein belonging to the lg superfamily. Fc $\epsilon$ Rl $\alpha$  forms a

tetrameric complex with one  $\beta$  and two  $\gamma$ -subunits. The Fc $\epsilon$ RI complex plays an important role in triggering IgE-mediated allergic reactions. It is abundantly expressed on mast and basophils and up-regulated by the presence of IgE. Following stimulation via Fc $\epsilon$ RI $\alpha$ , mast cells and basophils release bioactive chemical mediators such as histamine, resulting in the initiation of allergic reactions. Cross linking of the high-affinity receptor for IgE on tissue mast cells triggers immediate hypersensitivity with local symptoms. The MAR-1

monoclonal antibody reacts with the FcεRlα subunit.


Form: Liquid

Conjugation: FITC

Size: 50 Tests, 100 Tests, 200 Tests

Host Species: Armenian Hamster

**Isotype:** Armenian Hamster IgG



Ex:490 nm; Em:530 nm

**Isotype Control:** FITC Armenian Hamster IgG Isotype Control[PIP] [Product AGEL2314]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.