

Product Datasheet **PE Anti-Mouse FcεRlα Antibody [MAR-1]** Catalogue Code: AGEL2315

## Antibody Data

Applications: ECM	Product SKU:	AGEL2315	Clone:	MAR-1
	Applications:	FCM		
Reactivity: Mouse	Reactivity:	Mouse		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	High affinity immunoglobulin alpha;FcERI;Fcer1a;	epsilon receptor subunit alpha;Fc-epsilon RI-	
Uniprot ID:	P20489		
Background:	FccRl $\alpha$ is a transmembrane protein belonging to the Ig superfamily. FccRl $\alpha$ forms a tetrameric complex with one $\beta$ and two $\gamma$ -subunits. The FccRI complex plays an important role in triggering IgE-mediated allergic reactions. It is abundantly expressed on mast and basophils and up-regulated by the presence of IgE. Following stimulation via FccRI $\alpha$ , mast cells and basophils release bioactive chemical mediators such as histamine, resulting in the initiation of allergic reactions. Cross linking of the high-affinity receptor for IgE on tissue mast cells triggers immediate hypersensitivity with local symptoms. The MAR-1 monoclonal antibody reacts with the FccRI $\alpha$ subunit.		
Form:	Liquid	PE Excitation and Emission Spectra	
Conjugation:	PE	100 -	
Size:	50 Tests, 100 Tests, 200 Tests		
Host Species:	Armenian Hamster	80 - 00 - 00 - 00 - 00 - 00 - 00 - 00 -	
Isotype:	Armenian Hamster IgG	20	



Isotype Control: PE Armenian Hamster IgG Isotype Control[PIP] [Product AGEL2315]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.