

Product Datasheet **APC Anti-Mouse CD105 Antibody** [MJ7/18] Catalogue Code: AGEL2472

Antibody Data

Product SKU:	AGEL2472	Clone:	MJ7/18
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Endoglin;END;Eng;			
Uniprot ID:	Q63961			
Background:	CD105 is a 90 kD homodimeric type I integral membrane glycoprotein, also known as endoglin. It is expressed on endothelial cells (especially on angiogenic endothelial cells) and upregulated by hypoxia, activated monocytes, macrophages, bone marrow stromal cells, and some cytotrophoblasts. CD105 is a receptor for TGF- β 1, TGF- β 3 and modulates TGF- β signaling by interacting with TGF- β receptors I and/or II. CD105 also binds other growth factors such as actvin A, BMP-2, and BMP-7. CD105 has been show to be a useful marker for identifying proliferating endothelium involved in tumor angiogenesis and can be used for tumor imaging and prognosis, and has therapeutic potential for some solid tumors and other angiogenic diseases.			
Form:	Liquid	APC Excitation and Emission Spectra		
Conjugation:	APC	80		
Size:	50 Tests, 100 Tests, 200 Tests	(%) ps 60 -		
Host Species:	Rat	40 × 10		
Isotype:	Rat IgG2a, κ	20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)		
		Ex:650 nm; Em:660 nm		
Isotype Control:	APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL2472]			

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.