

#### **Product Datasheet**

# APC Anti-Mouse CD105 Antibody [MJ7/18]

Catalogue Code: AGEL2474

## Antibody Data

Product SKU: AGEL2474 Clone: MJ7/18

Applications: FCM

Reactivity: Mouse

### **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Endoglin; END; Eng;

Uniprot ID: Q63961

Background: CD105 is a 90 kD homodimeric type I integral membrane glycoprotein, also known as

endoglin. It is expressed on endothelial cells (especially on angiogenic endothelial cells) and upregulated by hypoxia, activated monocytes, macrophages, bone marrow stromal cells, and some cytotrophoblasts. CD105 is a receptor for TGF- $\beta$ 1, TGF- $\beta$ 3 and modulates TGF- $\beta$  signaling by interacting with TGF- $\beta$  receptors I and/or II. CD105 also binds other growth factors such as actvin A, BMP-2, and BMP-7. CD105 has been show to be a useful marker for identifying proliferating endothelium involved in tumor angiogenesis and can be used for tumor imaging and prognosis, and has therapeutic potential for some solid tumors

and other angiogenic diseases.


Form: Liquid

Conjugation: APC

Size: 25µg, 100µg

Host Species: Rat

**Isotype:** Rat IgG2a, κ



Ex:650 nm; Em:660 nm

**Isotype Control:** APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL2474]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/106 cells in 100  $\mu$ L

volume].