

Product Datasheet **FITC Anti-Human CD54 Antibody [15.2]** Catalogue Code: AGEL2808

Antibody Data

Product SKU:	AGEL2808	Clone:	15.2
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	Intercellular adhesion molecule 1;Icam1;MALA-2;MyD10;CD54;Icam-1; P05362 CD54 is a 85-110 kD type I transmembrane protein also known as ICAM-1. It is expressed on activated endothelial cells, high endothelial venules, T and B cells, monocytes/macrophages, granulocytes, and dendritic cells. The expression of ICAM-1 can be released from the cell surface. CD54 plays a role in cellular adhesion and is involved in inflammation and leukocyte extravasation. CD54 has also been shown to be the major cellular receptor for rhinovirus. ICAM-1 binds to CD11a/CD18 (LFA-1), CD11b/CD18 (Mac-1), CD11c/CD18 (p150, 95) as well as hyaluronan and fibrinogen.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	60 (S)	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 550 550 600 650 700 Wavelength (nm) Ex:490 nm; Em:530 nm	

Isotype Control: FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL2808]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.