

Product Datasheet

GenieFluor Red 780 Anti-Rat CD45 Antibody [OX-1]

Catalogue Code: AGEL3016

Antibody Data

Product SKU: AGEL3016 Clone: OX-1

Applications: FCM

Reactivity: Rat

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Receptor-type tyrosine-protein phosphatase C;Leukocyte common antigen; LCA; T200;

Ly-5;Ptprc;

Uniprot ID: P04157

Background: CD45 is a 180-220 kD protein also known as leukocyte common antigen (LCA). It is a

protein tyrosine phosphatase with multiple isoforms differing as a result of alternative splicing of the extracellular domain and glycosylation. CD45 is expressed on all hematopoietic cells except erythrocytes and platelets; isoform expression depends on cell type, activation state, and cell maturation. CD45 functions in signal transduction through T and B cell antigen receptors. CD45 has been shown to interact with various proteins including galectin-1, CD2, CD3, and CD4. The OX-1 antibody has been shown to partially

inhibit NK cell-mediated lysis of syngeneic tumor cells in vitro.


Form: Liquid

Conjugation: Genie FluorRed 780

Size: 25µg, 100µg

Host Species: Mouse

Isotype: Mouse IgG1, κ



Isotype Control: Genie Fluor Red 780 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3016]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/106 cells in 100 μ L

volume].