

#### **Product Datasheet**

# PE/Cyanine7 Anti-Human/Mouse KLRG-1 Antibody [2F1]

Catalogue Code: AGEL3110

### Antibody Data

Product SKU: AGEL3110 Clone: 2F1

Applications: FCM

Reactivity: Human; Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information:**

Alternate Names: MAFA; 2F1-Ag; Uniprot ID: Q96E93 O88713

Background: Killer cell lectin-like receptor G1 (KLRG1) is the mouse homolog of the rat mast cell

function-associated antigen (MAFA or 2F1-Ag). KLRG1 is a type II membrane glycoprotein that was first identified on the surface of rat mast cell line RBL-2H3. It is composed of a homodimer of glycosylated 30-38 kD subunits. Mouse and human homologs of KLRG1 are expressed by subsets of NK cells and lymphokine-activated killer (LAK) cells but not mast cells. KLRG1 is also expressed on subsets of CD8+ and CD4+ cells, including CD4+ and CD8+ effector/memory cells, potent regulatory CD4+ T cells. KLRG1 may be involved in regulating NK cell homeostasis. KLRG9 was found to recognize cadherins and thus inhibit immune responses by regulating the effector function and the developmental

processes of NK and T cells.


Form: Liquid

**Conjugation:** PE/Cyanine 7

Size: 20 Tests, 100 Tests, 200 Tests

**Host Species:** Syrian Hamster

**Isotype:** Syrian Hamster IgG



Ex:495;565;755 nm; Em:775 nm

Isotype Control: -

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.