

**PE/GenieFluor 594 Anti-Mouse  
CD45R/B220 Antibody [RA3.3A 1/6.1]**  
Catalogue Code: AGEL3253

## Antibody Data

Product SKU:	AGEL3253	Clone:	RA3.3A 1/6.1
Applications:	FCM		
Reactivity:	Mouse		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

**Alternate Names:** B220;

**Uniprot ID:** -

**Background:** CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-Fas<sup>lpr</sup> and MRL-Fas<sup>gld</sup> mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity.


**Form:** Liquid

**Conjugation:** PE/Genie Fluor594

**Size:** 50 Tests, 100 Tests, 200 Tests

**Host Species:** Rat

**Isotype:** Rat IgM,  $\kappa$



**Isotype Control:** PE/Genie Fluor 594 Rat IgM,  $\kappa$  Isotype Control[RTK2118] [Product AGEL3253]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

---