

Product Datasheet **FITC Anti-Human CD56 Antibody [MY31]** Catalogue Code: AGEL3297

Antibody Data

Product SKU:	AGEL3297	Clone:	MY31
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	Neural cell adhesion molecule 1;Leu-19;NKH1;NCAM-1;NCAM1;NCAM; P13591 CD56 is a single transmembrane glycoprotein also known as NCAM (neural cell adhesion		
Background.	molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is expressed on NK and NKT cells. CD56 is also expressed in the brain (cerebellum and cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparan sulfate.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	100	
Size:	20 Tests, 100 Tests, 200 Tests	g 60 -	
Host Species:	Mouse	60 40 -	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 550 550 600 650 700 Wavelength (nm) Ex:490 nm; Em:530 nm	

Isotype Control: FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL3297]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.