

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL3347</b>	<b>Clone:</b>	<b>RMV-7</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

<b>Alternate Names:</b>	Integrin alpha-V; $\alpha$ V integrin;Vitronectin Receptor;Integrin $\alpha$ V chain;ITGAV;
<b>Uniprot ID:</b>	P43406
<b>Background:</b>	CD51 is a 140 kD protein, also known as $\alpha$ V integrin, vitronectin receptor, and integrin $\alpha$ V. It is a member of the integrin family, expressed on activated T cells, polymorphonuclear granulocytes, platelets, blastocysts, and osteoclasts. CD51 forms heterodimers by association with integrins $\beta$ 1, $\beta$ 3, $\beta$ 5 or $\beta$ 6; these complexes then act as receptors for multiple extracellular matrix proteins (ECM). The $\alpha$ vintegrin heterodimers have varied functions in development, stimulation/activation and homeostasis. The primary ligands for CD51 complexes are fibronectin, fibrinogen, vitronectin, thrombospondin, von Willebrand factor, and CD31. The RMV-7 antibody has been reported to block binding of CD51 to vitronectin, fibronectin, and CD31 in some cell types, as well as blocking LAK cell cytotoxicity.
<b>Form:</b>	Liquid
<b>Conjugation:</b>	Biotin
<b>Size:</b>	25&micro;g, 100&micro;g
<b>Host Species:</b>	Rat
<b>Isotype:</b>	Rat IgG1, $\kappa$
<b>Isotype Control:</b>	Biotin Rat IgG1, $\kappa$ Isotype Control[HRPN] [Product AGEL3347]
<b>Storage Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \mu\text{g}$  per  $10^6$  cells in 100  $\mu\text{L}$  volume or 100  $\mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

---