

Product Datasheet **APC Anti-Mouse CD163 Antibody [S15049F]** Catalogue Code: AGEL3395

Antibody Data

Product SKU:	AGEL3395	Clone:	S15049F
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	M130; MM130; SCARI1;		
Uniprot ID:	Q2VLH6		
Background:	CD163 is a member of the group B scavenger receptor cysteine-rich superfamily, also known as GHI/61, M130, RM3/1, p155, hemoglobin-haptoglobin complex receptor, or macrophage-associated antigen. It is a 134 kD (non-reduced)/155 kD (reduced) glycoprotein primarily expressed on macrophages, Kupffer cells, monocytes, a subset of dendritic cells, and a subset of hematopoietic stem/progenitor cells. CD163 binds to haptoglobin-hemoglobin complex and TWEAK, and plays a role in clearing hemoglobin and regulating cytokine production by macrophages. Membrane CD163 can be cleaved by metalloproteinases (MMP), resulting in a soluble form. Elevated serum level of sCD163 has been implicated in many kinds of inflammatory diseases.		
Form:	Liquid	APC Excitation and Emission Spectra	
Conjugation:	APC	80 -	
Size:	50 Tests, 100 Tests, 200 Tests		
Host Species:	Rat	Normalized (%)	
Isotype:	Rat IgG2a, к	20	
		350 480 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:650 nm; Em:660 nm	
Isotype Control:	APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL3395]		
Storage Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.