

Product Datasheet **FITC Anti-Human CD62L Antibody [HI62L]** Catalogue Code: AGEL3484

Antibody Data

Product SKU:	AGEL3484	Clone:	HI62L
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	L-selectin;Sell;CD62 antigen-like family member L;LAM-1;LECAM1;Lymph node homing receptor;Ly-22; CD62L;Lnhr;Ly22;		
Uniprot ID:	P14151		
Background:	CD62L is a 74-95 kD single chain type I glycoprotein referred to as L-selectin or LECAM- 1. It is expressed on most peripheral blood B cells, subsets of T and NK cells, monocytes, granulocytes, and certain hematopoietic malignant cells. CD62L binds to carbohydrates present on certain glycoforms of CD34, glycam-1, and MAdCAM-1 and with a low affinity to anionic oligosaccharide sequences related to sialylated Lewis X (sLex, CD15s) through its C-type lectin domain. CD62L is important for the homing of naïve lymphocytes to high endothelial venules in peripheral lymph nodes and Peyer's patches. It also plays a role in leukocyte rolling on activated endothelial cells.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	60 - 40	
Isotype:	Mouse IgG2a, κ		
		350 400 450 500 550 600 650 700 Wavelength (nm)	
		Ex:490 nm; Em:530 nm	
Isotype Control:	EITC Mouse IgG2a, K Isotype Control[C1 18 4] [Product AGEL 3484]		

Isotype Control: FITC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL3484]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.