



## Product Datasheet

### Purified Anti-Mouse IL-17A Antibody [17F3]

Catalogue Code: AGEL3512

#### Antibody Data

<b>Product SKU:</b>	<b>AGEL3512</b>	<b>Clone:</b>	<b>17F3</b>
<b>Applications:</b>	<b>ICFCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

#### Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

#### Product Information:

<b>Alternate Names:</b>	Interleukin-17A;IL17a;Ctla9;
<b>Uniprot ID:</b>	Q62386
<b>Background:</b>	The 17F3 monoclonal antibody reacts with mouse IL-17A a 15-20 kDa cytokine expressed by Th17 cells, $\gamma\delta$ T cells, iNKT cells, NK cells, LTI cells, neutrophils, and intestinal Paneth cells. IL-17A has pleiotropic effects in immunoregulation and inflammation. It plays an important role in anti-microbial and chronic inflammation by inducing cytokine and chemokine production, neutrophil influx, and the production of antibacterial peptides but it is also an inflammatory mediator in the development of autoimmune diseases including rheumatoid arthritis, asthma, multiple sclerosis, and psoriasis. The 17F3 antibody has been shown to neutralize IL-18A in vivo.
<b>Form:</b>	Liquid
<b>Conjugation:</b>	Unconjugated
<b>Size:</b>	25&micro;g, 100&micro;g
<b>Host Species:</b>	Mouse
<b>Isotype:</b>	Mouse IgG1, $\kappa$
<b>Isotype Control:</b>	Purified Mouse IgG1, $\kappa$ Isotype Control[MOPC-21] [Product AGEL3512]
<b>Storage Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \mu\text{g}$  per  $10^6$  cells in 100  $\mu\text{L}$  volume or 100  $\mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

---