## CAB0440



## **Product Information**

| Product SKU: | CAB0440 | Gene ID:      | 4087   | Size:       | 20uL, 100uL  |
|--------------|---------|---------------|--------|-------------|--------------|
| Clone No:    | -       | Host Species: | Rabbit | Reactivity: | Human, Mouse |
|              |         |               |        |             |              |

## **Additional Information**

| Observed MW:   | 50kDa | Conjugate: | Unconjugated |
|----------------|-------|------------|--------------|
| Calculated MW: | 52kDa | lsotype:   | lgG          |

## **Immunogen Information**

| Background            | The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products   |
|-----------------------|---|
|                       | of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD          |
|                       | proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways.  |
|                       | This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates         |
|                       | multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is  |
|                       | recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation  |
|                       | (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. |
|                       | The phosphorylation induces the dissociation of this protein with SARA and the association with the       |
|                       | family member SMAD4. The association with SMAD4 is important for the translocation of this protein        |
|                       | into the nucleus, where it binds to target promoters and forms a transcription repressor complex with     |
|                       | other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates  |
|                       | the signal from the activin. Alternatively spliced transcript variants have been observed for this gene.  |
| Recommended Dilution: | WB,1:500 - 1:1000 IHC-P,1:50 - 1:200  |
| Synonyms:             | JV18; LDS6; CHTD8; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2; Smad2  |
| Purifcation Method:   | Affinity purification   |
| Immunogen:            | Recombinant fusion protein containing a sequence corresponding to amino acids 175-270 of human            |
|                       | Smad2 (NP_005892.1).  |
| Storage:              | Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.          |