## **ICAM-1 Mouse Monoclonal Antibody**



## **CAB11869**

**Product Information** 

**Protein Background** 

Size:

This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18

and is also exploited by Rhinovirus as a receptor.

50uL, 100uL

**Observed MW:** 

Immunogen information

95kDa

Gene ID:

Calculated MW: 3383

57kDa

Uniprot

**Applications:** P05362

WB IF

Synonyms:

**Reactivity:** BB2; CD54; P3.58; ICAM1

Human

Immunogen:

Antibody Information Recombinant protein of human ICAM1

**Recommended dilutions:** 

WB 1:500 - 1:2000 IF 1:50 -

1:200

Source:

Mouse

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.


Isotype:

IgG


**Purification:** 

Affinity purification


## **Product Images**



Western blot analysis of extracts of Raji cells, using ICAM1 antibody (CAB11869). Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (CABS003) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of extracts of various cell lines, using ICAM1 antibody (CAB11869). Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (CABS003) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of SW480 cells using ICAM1 antibody (CAB11869). Blue: DAPI for nuclear staining.