

## CAB21442

## **Product Information**

Product SKU:	CAB21442	Gene ID:	2908	Size:	20uL, 100uL
Clone No:	-	Host Species:	Rabbit	Reactivity:	Human,Rat

## **Additional Information**

Observed MW:	Refer to figures	Conjugate:	Unconjugated
Calculated MW:	86kDa	lsotype:	lgG

## **Immunogen Information**

Background:This gene encodes glucocorticold receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GCR; GCRST; Glucocorticoid ReceptorPurification Method:Affinity purification Glucocorticoid Receptor (NP_001191194.1).Storage:Storage:Storage / tar -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.	De al-ana an di	This same encodes all second and the birds are firstly both as a transmistion forter that his de
Heir transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GCL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purification Glucocorticoid Receptor (NP_001191194.1).	Background:	This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds
the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate
<ul> <li>responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).</li> <li>Recommended Dilution: WB,1:500 - 1:2000 IHC-P,1:50 - 1:200</li> <li>Synonyms: GR; GCR; GRL; GCCR; GCRST; Glucocorticoid Receptor</li> <li>Purifcation Method: Affinity purification</li> <li>Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).</li> </ul>		their transcription, and as a regulator of other transcription factors. This receptor is typically found in
associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory
transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended DilutionWB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are
the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in
Immunogen:functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).Recommended Dilution:WB 1:500 - 1:2000 IHC-P.1:50 - 1:200Synonyms:GR; GCR; GCR; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human glucocorticoid Receptor (NP_001191194.1).		transcript variants encoding either the same or different isoforms. Additional isoforms resulting from
activities (PMID:15866175).Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		the use of alternate in-frame translation initiation sites have also been described, and shown to be
Recommended Dilution:WB,1:500 - 1:2000 IHC-P,1:50 - 1:200Synonyms:GR; GCR; GCR; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional
Synonyms:GR; GCR; GRL; GCCR; GCRST; Glucocorticoid ReceptorPurifcation Method:Affinity purificationImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).		activities (PMID:15866175).
Purifcation Method:       Affinity purification         Immunogen:       Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human Glucocorticoid Receptor (NP_001191194.1).	<b>Recommended Dilution</b> :	WB,1:500 - 1:2000 IHC-P,1:50 - 1:200
Immunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of humanGlucocorticoid Receptor (NP_001191194.1).	Synonyms:	GR; GCR; GRL; GCCR; GCRST; Glucocorticoid Receptor
Glucocorticoid Receptor (NP_001191194.1).	Purifcation Method:	Affinity purification
	Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human
<b>Storage</b> : Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.		Glucocorticoid Receptor (NP_001191194.1).
	Storage:	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.