

## **Product Information**

| Product SKU: | CAB21574 | Gene ID:      | 5696   | Size:               | 20uL, 100uL       |
|--------------|----------|---------------|--------|---------------------|-------------------|
| Clone No:    | -        | Host Species: | Rabbit | <b>Reactivity</b> : | Human, Mouse, Rat |
|              |          |               |        |                     |                   |

## **Additional Information**

| Observed MW:   | Refer to figures | Conjugate: | Unconjugated |
|----------------|------------------|------------|--------------|
| Calculated MW: | 30kDa            | lsotype:   | lgG          |

## **Immunogen Information**

| Background                    | The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core         |  |  |  |
|-------------------------------|--|--|--|--|
|                               | structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed  |  |  |  |
|                               | of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed             |  |  |  |
|                               | throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent    |  |  |  |
|                               | process in a non-lysosomal pathway. An essential function of a modified proteasome, the                  |  |  |  |
|                               | immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the           |  |  |  |
|                               | proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is    |  |  |  |
|                               | located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is |  |  |  |
|                               | induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5        |  |  |  |
|                               | subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit.       |  |  |  |
|                               | Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to   |  |  |  |
|                               | yield the same mature subunit.   |  |  |  |
| <b>Recommended Dilution</b> : | WB,1:500 - 1:2000 IHC-P,1:50 - 1:200 IF/ICC,1:50 - 1:200   |  |  |  |
| Synonyms:                     | JMP; ALDD; LMP7; NKJO; D6S216; PRAAS1; PSMB5i; RING10; D6S216E; PSMB8                                    |  |  |  |
| Purifcation Method:           | Affinity purification  |  |  |  |
| Immunogen:                    | Recombinant fusion protein containing a sequence corresponding to amino acids 163-272 of human           |  |  |  |
|                               | PSMB8 (NP_004150.1).   |  |  |  |
| Storage:                      | Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.       |  |  |  |