## **Cyclin H Monoclonal Antibody**





**Product Information** 

**Product SKU**: CAB21885 **Gene ID**: 902 **Size**: 20uL, 100uL

Clone No: ARC2823 Host Species: Rabbit Reactivity: Human, Mouse

**Additional Information** 

**Observed MW**: 35kDa **Conjugate:** Unconjugated

Calculated MW: 38kDa Isotype: IgG

Immunogen Information

**Background**: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are

characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control

and the cell cycle machinery. A pseudogene of this gene is found on chromosome 4. Alternate splicing

results in multiple transcript variants.

**Recommended Dilution**: WB,1:500 - 1:1000

**Synonyms**: CAK; p34; p37; CycH; Cyclin H

**Purifcation Method**: Affinity purification

**Immunogen**: A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Cyclin H (P51946).

**Storage**: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,0.05% BSA,50%

glycerol,pH7.3.