

CAB22002

Product Information

Product SKU:	CAB22002	Gene ID:	5045	Size:	20uL, 100uL
Clone No:	-	Host Species:	Rabbit	Reactivity:	Human

Additional Information

Observed MW:	90kDa	Conjugate:	Unconjugated
Calculated MW:	87kDa	Isotype:	IgG

Immunogen Information

Background: This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. It encodes a type 1 membrane bound protease that is expressed in many tissues, including neuroendocrine, liver, gut, and brain. The encoded protein undergoes an initial autocatalytic processing event in the ER and then sorts to the trans-Golgi network through endosomes where a second autocatalytic event takes place and the catalytic activity is acquired. Like other members of this convertase family, the product of this gene specifically cleaves substrates at single or paired basic residues. Some of its substrates include parathyroid hormone, transforming growth factor beta 1 precursor, proalbumin, pro-beta-secretase, membrane type-1 matrix metalloproteinase, beta subunit of pro-nerve growth factor and von Willebrand factor. It is thought to be one of the proteases responsible for the activation of HIV envelope glycoproteins gp160 and gp140, and may play a role in tumor progression. Unlike SARS-CoV and other coronaviruses, the spike protein of SARS-CoV-2 is thought to be uniquely cleaved by this protease. Alternative splicing results in multiple transcript variants.

Recommended Dilution: WB,1:500 - 1:1000

Synonyms: FUR; PACE; SPC1; PCSK3; Furin

Purification Method: Affinity purification

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 200-300 of human Furin (NP_002560.1).

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.