

CAB22368

Product Information								
Product SKU:	CAB22368	Gene ID:	1489668		Size:	20uL, 100uL		
Clone No:	ARC52570	Host Species:	Rabbit		Reactivity :	SARS-CoV		
Additional Information								
Observed MW:	181KD		Conjugate:	Unconjugate	d			
Calculated MW	: 139kDa		lsotype:	IgG				

Immunogen Information

Background:	The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors
	on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl
	peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion
	molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity
	and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins
	on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be
	essential in the binding of the virus to the host cell at the advent of the infection process. It's been
	reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory
	epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I
	transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding
	domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements
	needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody
	and T-cell responses, as well as protective immunity. The main functions for the Spike protein are
	summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and
	specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine
	design; Can be transmitted between different hosts through gene recombination or mutation of the
	receptor binding domain (RBD), leading to a higher mortality rate.
Recommended Dilution:	WB,1:1000 - 1:5000
Synonyms:	E2; SARS-CoV Spike RBD
Purifcation Method:	Affinity purification
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 350-450 of coronavirus SARS-CoV
	Spike RBD (NP_828851.1).

Contact Details | Dublin, Ireland Email: techsupport@assaygenie.com | Web: <u>www.assaygenie.com</u> Copyright © 2020 Reagent Genie, All Rights Reserved. All information / detail is correct at time of going to print.

Storage:

Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.