## CAB22570



## **Product Information**

Product SKU:	CAB22570	Gene ID:	3574		Size:	20uL, 100uL
Clone No:	ARC57489	Host Species:	Rabbit		<b>Reactivity</b> :	Human
Additional Ir	nformation					
Observed MW:	30-35kDa		Conjugate:	Unconjugated	I	

Isotype:

lgG

## **Immunogen Information**

20kDa

Calculated MW:

Background	The protein encoded by this gene is a cytokine important for B and T cell development. This cytokine
	and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-
	stimulating factor. IL7 is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB)
	during early T cell development. This cytokine can be produced locally by intestinal epithelial and
	epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. IL7 plays
	an essential role in lymphoid cell survival, and in the maintenance of naive and memory T cells.
	Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice
	variants have been described but their presence in normal tissues has not been confirmed. Severe acute
	respiratory syndrome coronavirus 2 (SARS-CoV-2) infection can be a potent inducer of proinflammatory
	cytokines and chemokines which may defend against the infection, but may also mediate destructive
	lung injury. Elevated serum IL7 levels, together with several other circulating cytokines and chemokines,
	has been found to be associated with the severity of Coronavirus Disease 19 (COVID-19).
Recommended Dilution:	WB,1:10000 - 1:30000
Synonyms:	IL-7; IL7
Purifcation Method:	Affinity purification
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 26-177 of human IL7
	(NP_000871.1).
Storage:	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,0.05% BSA,50%
	glycerol,pH7.3.