

## CAB4998

---

### Product Information

<b>Product SKU:</b>	CAB4998	<b>Gene ID:</b>	5581	<b>Size:</b>	20uL, 100uL
<b>Clone No:</b>	ARC1270	<b>Host Species:</b>	Rabbit	<b>Reactivity:</b>	Mouse

---

### Additional Information

<b>Observed MW:</b>	82kDa	<b>Conjugate:</b>	Unconjugated
<b>Calculated MW:</b>	84kDa	<b>Isotype:</b>	IgG

---

### Immunogen Information

**Background:** Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been shown to be involved in many different cellular functions, such as neuron channel activation, apoptosis, cardioprotection from ischemia, heat shock response, as well as insulin exocytosis. Knockout studies in mice suggest that this kinase is important for lipopolysaccharide (LPS)-mediated signaling in activated macrophages and may also play a role in controlling anxiety-like behavior.

**Recommended Dilution:** WB, 1:500 - 1:2000

**Synonyms:** PKCE; nPKC-epsilon; PKC epsilon

**Purification Method:** Affinity purification

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 500-600 of human PKC epsilon (Q02156).

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH 7.3.