Phospho-Smad2-S465/S467 + Smad3-S423/S425 Monoclonal Antibody



CABP1343

Product Information

Product SKU: CABP1343 **Gene ID**: 4087/4088 **Size**: 20uL, 100uL

Clone No: ARC56314 Host Species: Rabbit Reactivity: Human, Mouse, Rat

Additional Information

Observed MW: 52kDa/60kDa **Conjugate:** Unconjugated

Calculated MW: 48kDa/52kDa/25kDa/35kDa/43 Isotype: IgG

kDa

Immunogen Information

Background: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products

of the Drosophila gene & apos; mothers against decapentaplegic & apos; (Mad) and the C. elegans gene

Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple

signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and

thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This

protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor

activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-

beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the

association with the family member SMAD4. The association with SMAD4 is important for the

translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin

type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants

have been observed for this gene. [provided by RefSeq, May 2012]

Recommended Dilution: WB,1:1000 - 1:5000

Synonyms: HSPC193; HsT17436; JV15-2; LDS1C; LDS3; MADH3; Phospho-Smad2-S465/S467 + Smad3-S423/S425

Purifcation Method: Affinity purification

Immunogen: A synthetic phosphorylated peptide around S465 & S467 of human SMAD2 (NP_005892.1).

Storage: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300,0.05% BSA,50%

glycerol,pH7.3.