

# Anti-CD43 Recombinant Rabbit Monoclonal Antibody

## Product Information

**Product SKU:**

HDAB0202

**Size:**

100 µg

**Isotype:**

Rabbit IgG

**Host:**

Rabbit

**Clonality:**

Monoclonal

**Clone:**

DM211

**Category:**

Recombinant Antibody

**Reactivity:**

Human

**Synonyms:**

CD43, GALGP, GPL115, LSN

**Formulation:**

Powder

**Purification Method:**

Purified from cell culture supernatant by affinity chromatography

**Buffer:**

1XPBS

**Storage and Stability:**

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

**Applications**

ELISA, Flow Cyt

**Recommended Dilutions:**

ELISA 1:5000-10000; Flow Cyt 1:100

**Background:**

This gene encodes a highly sialylated glycoprotein that functions in antigen-specific activation of T cells, and is found on the surface of thymocytes, T lymphocytes, monocytes, granulocytes, and some B lymphocytes. It contains a mucin-like extracellular domain, a transmembrane region and a carboxy-terminal intracellular region. The extracellular domain has a high proportion of serine and threonine residues, allowing extensive O-glycosylation, and has one potential N-glycosylation site, while the carboxy-terminal region has potential phosphorylation sites that may mediate transduction of activation signals. Different glycoforms of this protein have been described. In stimulated immune cells, proteolytic cleavage of the extracellular domain occurs in some cell types, releasing a soluble extracellular fragment. Defects in expression of this gene are associated with Wiskott-Aldrich syndrome. [provided by RefSeq, Sep 2017]

**Research Use Only:**

Research Use Only