

# Anti-IL5 Chimeric Recombinant Rabbit Monoclonal Antibody

## Product Information

**Product SKU:**

HDAB0232

**Size:**

100 µg

**Isotype:**

Rabbit/Human Fc chimeric IgG1

**Host:**

Rabbit

**Clonality:**

Monoclonal

**Clone:**

DMC274

**Category:**

Recombinant Antibody

**Reactivity:**

Human

**Synonyms:**

IL-5, TRF, Interleukin-5, EDF

**Formulation:**

Powder

**Purification Method:**

Purified from cell culture supernatant by affinity chromatography

**Buffer:**

1XPBS

**Storage and Stability:**

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

**Applications**

Flow Cyt

**Recommended Dilutions:**

Flow Cyt 1:100

**Background:**

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukin 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2 . This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31.

**Research Use Only:**

Research Use Only