

Anti-VEGFA Chimeric Recombinant Rabbit Monoclonal Antibody

Product Information

Product SKU:

HDAB0234

Size:

100 µg

Isotype:

Rabbit/Human Fc chimeric IgG1

Host:

Rabbit

Clonality:

Monoclonal

Clone:

DMC276

Category:

Recombinant Antibody

Reactivity:

Human

Synonyms:

MVCD1, VEGF, VPF

Formulation:

Powder

Purification Method:

Purified from cell culture supernatant by affinity chromatography

Buffer:

1XPBS

Storage and Stability:

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Applications

Flow Cyt

Recommended Dilutions:

Flow Cyt 1:100

Background:

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.

Research Use Only:

Research Use Only