IVMB0520



Product Information Product SKU: IVMB0520 Clone: GSK2857914 Target: **BCMA** Size: 500 µg Isotype: Human IgG1ĸ Additional Information **Reactivity:** Human Host Species: Human Antibody Type: Biosimilar Recombinant Human Monoclonal Antibody Expression Host: HEK-293 Cells

Immunogen Information

Background:B cell maturation antigen (BCMA, CD269, TNFRSF-17) is a type III transmembrane
glycoprotein that is a member of the tumor necrosis factor (TNF) receptor superfamily¹.
BCMA functions as a cell-surface receptor and is involved in the regulation of B cell
proliferation, maturation, and differentiation into plasma cells, and is also required for the
survival of long-lived plasma cells¹, ². BCMA is more abundantly expressed on malignant
plasma cells than normal plasma cells and is a novel treatment target for multiple myeloma
(MM)¹, ³, ⁴, a plasma cell malignancy characterized by clonal proliferation of plasma cells
within the bone marrow².

BCMA expression is upregulated during MM pathogenesis and evolution, with higher levels associated with poorer prognosis¹. The soluble form of BCMA, which is derived from direct shredding of membrane BMCA through γ -secretase activity, is also significantly elevated in MM patients relative to healthy individuals and is associated with worse clinical responses.

Belantamab (J6M0) is a novel afucosylated, humanized antagonistic anti-BCMA IgG1 monoclonal antibody⁴ produced in a Chinese Hamster Ovary cell line using recombinant DNA technology⁵. Belantamab has been used in clinical trials as part of the antibody conjugate belantamab mafodotin-blmf (GSK2857916)⁵, ⁶, ⁷ and has been shown to directly and indirectly target MM cells via multiple mechanisms of action⁴. Binding is BCMA-specific, with belantamab competing with BCMA's two ligands BAFF and APRIL and also inhibiting



ligand-induced NF κ B signaling⁴. The afucosylation significantly increases the binding affinity of the Fc domain to the Fc γ R (Fc γ RIIIa) expressed on effector cells and enhances antibody-dependent cell-mediated cytotoxicity (ADCC)⁴.

Endotoxin Level: < 1.0 EU/mg as determined by the LAL method

Applications: ELISA

Synonyms: GSK2857914, TNFRSF17, BCMA, CD269

Antigen Distribution: BCMA protein is expressed on the surface of normal B lymphocytes and nearly all multiple myeloma cell lines. BCMA is almost exclusively expressed on plasmablasts and plasma cells and is also weakly expressed on some memory B cells committed to plasma cell differentiation and on plasmacytoid dendritic cells. BCMA is nearly absent on naïve and memory B cells.

Immunogen: Human TNFRSF17/CD269 (BCMA)

- Formulation:This biosimilar antibody is aseptically packaged and formulated in 0.01 M phosphate
buffered saline (150 mM NaCl) PBS pH 7.2 7.4 with no carrier protein, potassium, calcium
or preservatives added. Due to inherent biochemical properties of antibodies, certain
products may be prone to precipitation over time. Precipitation may be removed by aseptic
centrifugation and/or filtration.
- Specificity:This non-therapeutic biosimilar antibody uses the same variable region sequence as the
therapeutic antibody Belantamab. This product is for research use only. Belantamab activity
is directed against Human BCMA.
- Product Preparation:
 Recombinant biosimilar antibodies are manufactured in an animal free facility using onlyin vitroprotein free cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.
- Storage & Handling:Functional grade biosimilar antibodies may be stored sterile as received at 2-8°C for up to
one month. For longer term storage, aseptically aliquot in working volumes without diluting
and store at -80°C. Avoid Repeated Freeze Thaw Cycles.