

## Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse, Rat

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

ELISA:1:2000-1:5000, WB:1:500-1:2000,  
IHC:1:50-1:200

**Protein Background:**

The protein encoded by this gene belongs to the class-II aminoacyl-tRNA synthetase family. Aminoacyl-tRNA synthetases play critical roles in mRNA translation by charging tRNAs with their cognate amino acid. The encoded protein is a mitochondrial enzyme that specifically aminoacylates alanyl-tRNA. Mutations in this gene are a cause of combined oxidative phosphorylation deficiency 8. Catalyzes the attachment of alanine to tRNA(Ala) in a two-step reaction: alanine is first activated by ATP to form Ala-AMP and then transferred to the acceptor end of tRNA(Ala). Also edits incorrectly charged tRNA(Ala) via its editing domain.

**Gene ID:**

AARS2

**Uniprot**

Q5JZ9

**Synonyms:**

Alanyl-tRNA synthetase 2, mitochondrial

**Immunogen:**


Fusion protein of human AARS2.

**Storage:**


-20&deg; C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

## Product Images


---



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO13998(AARS2 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: x—200).



Gel: 8%SDS-PAGE, Lysate: 40 &mu; g, Lane: SP20 cells, Primary antibody: PACO13998(AARS2 Antibody) at dilution 1/600, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 15 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human lymphoma tissue using PACO13998(AARS2 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: x—200).