## **ATP5J Antibody**

# AssayGenie 🗳

#### PACO14120

#### **Product Information**

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

lgG

**Applications:** 

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:15-1:50

**Protein Background:** 

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the F6 subunit of the Fo complex, required for F1 and Fo interactions. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. A pseudogene exists on chromosome Yp11.

Gene ID:

ATP5J

Uniprot

P18859

**Synonyms:** 

ATP synthase, H+ transporting, mitochondrial Fo complex, subunit F6


Immunogen:


Fusion protein of human ATP5J.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


### **Product Images**





The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using PACO14120(ATP5J Antibody) at dilution 1/15, on the right is treated with fusion protein. (Original magnification: x—200).

Gel: 12%SDS-PAGE, Lysate: 40 μ g, Lane 1-4: Human fetal brain tissue, 293T cells, mouse intestinum tenue tissue, lovo cells, Primary antibody: PACO14120(ATP5J Antibody) at dilution 1/300, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute.



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO14120(ATP5J Antibody) at dilution 1/15, on the right is treated with fusion protein. (Original magnification: x—200).