## **ANKRD54 Antibody**



## PACO15447

Reactivity:

## **Product Information**

Size: Protein Background:

50ul Plays an important role in regulating intracellular signaling events associated with erythroid terminal differentiation. Interacts (via ankyrin repeat region) with LYN (via

SH3-domain) in an activation-independent status of LYN By similarity. Forms a

Human, Mouse, Rat multiprotein complex with LYN and HCLS1 By similarity. Interacts with TSN2, VAV1,

DBNL AND LASP1.

Source:

Gene ID:

Rabbit ANKRD54

Isotype: Uniprot

lgG Q6NXT1

Applications: Synonyms:

ELISA, WB, IHC ankyrin repeat domain 54


Recommended dilutions: Immunogen:


ELISA:1:2000-1:5000, WB:1:200-1:1000, IHC:1:50-1:200 Fusion protein of human ANKRD54.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


## **Product Images**





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO15447(ANKRD54 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: x—200).

Gel: 10%SDS-PAGE, Lysate: 40 μ g, Lane 1-2: Mouse testis tissue, human colon cancer tissue, Primary antibody: PACO15447(ANKRD54 Antibody) at dilution 1/250, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO15447(ANKRD54 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: x—200).